

SEARCH FOR NEUTRINO MASS AND DARK MATTER IN UNDERGROUND EXPERIMENTS

H.V. Klapdor-Kleingrothaus

Max-Planck-Institut für Kernphysik,

P.O. Box 10 39 80, D-69029 Heidelberg, Germany

Spokesman of HEIDELBERG-MOSCOW and GENIUS Collaborations

E-mail: klapdor@gustav.mpi-hd.mpg,

Home-page: http://www.mpi-hd.mpg.de/non_acc/

Search for the neutrino mass and for cold dark matter in the Universe are at present two of the most exiting fields of particle physics and cosmology. This lecture will restrict itself on the search for neutralinos as cold dark matter, and for the absolute scale of the masses of neutrinos, which are the favoured hot dark matter candidates.

1 Introduction

In this lecture we shall discuss two central problems of particle physics and cosmology. The neutrino mass is one of the key quantities in grand unified theories, and at the same time candidate for hot dark matter in the Universe. Supersymmetry is regarded as the most natural extension of present particle physics theories. The lightest SUSY particle, usually assumed to be the neutralino, is the favored candidate for cold dark matter. We shall discuss here the search for the neutrino mass, concentrating on double beta decay, and the search for neutralinos, in underground experiments.

The neutrino oscillation interpretation of the atmospheric and solar neutrino data, deliver a strong indication for a non-vanishing neutrino mass. While such kind of experiments yields information on the difference of squared neutrino mass eigenvalues and on mixing angles, they cannot fix the absolute scale of the neutrino mass. Information from double beta decay experiments is indispensable to solve this question^{9,10,11,12}. Another important problem is that of the fundamental character of the neutrino, whether it is a Dirac or a Majorana particle^{2,3}. Neutrinoless double beta decay could answer also this question. Perhaps the main question, which can be investigated by double beta decay with high sensitivity, is that of lepton number conservation or non-conservation.

The physics motivations to search for dark matter are manifold.

Recent investigation of the cosmic microwave background radiation (MAXIMA, BOOMERANG, DASI) together with large scale structure results fix $\Omega_\Lambda + \Omega_m \sim 1$, where Ω_Λ ($= \rho_\Lambda/\rho_c$) stands for dark energy and

<i>HVKK-Triest01-Prepr: submitted to World Scientific on February 7, 2008</i>
--

Ω_m for matter. With the early nucleosynthesis constraint of $\Omega_{bar} \sim 0.04$ the need for non-baryonic dark matter is evident. This is true even for our galaxy, since MACHOs represent only a small fraction of galactic dark matter⁶².

Although there exist other candidates such as axions⁶³, gravitinos, etc., neutralinos seem to be the favored candidates at present.

Hot dark matter, according to CMB and LSS (Redshift-Survey) results still contributes up to 38% of the dark matter⁸⁶. This corresponds to a sum of neutrino masses < 5.5 eV.

Our present picture of the mass/energy distribution in the Universe is as given in Fig.1.




Figure 1. Our present understanding of the mass composition of the Universe.

In section 2, we discuss double beta decay, including in section 2.2 the expectation from neutrino oscillation experiments, and in section 2.3 the recent evidence for the neutrinoless decay mode from the HEIDELBERG-MOSCOW experiment. In section 2.4 we discuss the future of the field.

In section 3 we shall discuss dark matter search. We start in 3.1 with the expectation from SUSY models and then discuss in section 3.2 the present status of cold dark matter experiments - including the important indication from the DAMA experiment - and its future. In section 3.3 we give some comments to neutrinos as hot dark matter. In section 4 we give a conclusion.

2 Neutrino Masses and Double Beta Decay

2.1 General

Double beta decay, the rarest known nuclear decay process, can occur in different modes:

$$2\nu\beta\beta - \text{decay} : \quad A(Z, N) \rightarrow A(Z+2, N-2) + 2e^- + 2\bar{\nu}_e \quad (1)$$

$$0\nu\beta\beta - \text{decay} : \quad A(Z, N) \rightarrow A(Z+2, N-2) + 2e^- \quad (2)$$

$$0\nu(2)\chi\beta\beta - \text{decay} : \quad A(Z, N) \rightarrow A(Z+2, N-2) + 2e^- + (2)\chi \quad (3)$$

While the two-neutrino mode (1) is allowed by the Standard Model of particle physics, the neutrinoless mode ($0\nu\beta\beta$) (2) requires violation of lepton number ($\Delta L=2$). This mode is possible only, if the neutrino is a Majorana particle, i.e. the neutrino is its own antiparticle (E. Majorana ², G. Racah ³, for subsequent works we refer to ^{5,6,7}, for some reviews see ^{14,15,16,17,12,18}). First calculations of $0\nu\beta\beta$ decay based on the Majorana theory have been done by W.H. Furry ⁴.

Neutrinoless double beta decay can not only probe a Majorana neutrino mass, but various new physics scenarios beyond the Standard Model, such as R-parity violating supersymmetric models, R-parity conserving SUSY models, leptoquarks, violation of Lorentz-invariance, and compositeness (for a review see ^{12,40,41}). Any theory containing lepton number violating interactions can in principle lead to this process allowing to obtain information on the specific underlying theory. The experimental signature of the neutrinoless mode is a peak at the Q-value of the decay.

The unique feature of neutrinoless double beta decay is that a measured half-life allows to deduce information on the effective Majorana neutrino mass $\langle m \rangle$, which is a superposition of neutrino mass eigenstates: ^{14,15}

$$\begin{aligned} [T_{1/2}^{0\nu}(0_i^+ \rightarrow 0_f^+)]^{-1} = & C_{mm} \frac{\langle m \rangle^2}{m_e^2} + C_{\eta\eta} \langle \eta \rangle^2 + C_{\lambda\lambda} \langle \lambda \rangle^2 + C_{m\eta} \langle \eta \rangle \frac{\langle m \rangle}{m_e} \\ & + C_{m\lambda} \langle \lambda \rangle \frac{\langle m \rangle}{m_e} + C_{\eta\lambda} \langle \eta \rangle \langle \lambda \rangle, \end{aligned} \quad (1)$$

$$\langle m \rangle = |m_{ee}^{(1)}| + e^{i\phi_2} |m_{ee}^{(2)}| + e^{i\phi_3} |m_{ee}^{(3)}|, \quad (2)$$

where $m_{ee}^{(i)} \equiv |m_{ee}^{(i)}| \exp(i\phi_i)$ ($i = 1, 2, 3$) are the contributions to $\langle m \rangle$ from individual mass eigenstates, with ϕ_i denoting relative Majorana phases connected with CP violation, and $C_{mm}, C_{\eta\eta}, \dots$ denote nuclear matrix elements,

which can be calculated, (see, e.g. ²³, for a review see e.g. ^{24,12,15,16,44}). Ignoring contributions from right-handed weak currents on the right-hand side of eq.(1), only the first term remains.

2.2 Allowed Ranges of $\langle m \rangle$ by ν Oscillation Experiments

The observable of double beta decay

$$\langle m \rangle = |m_{ee}^{(1)}| + e^{i\phi_2}|m_{ee}^{(2)}| + e^{i\phi_3}|m_{ee}^{(3)}|,$$

with U_{ei} denoting elements of the neutrino mixing matrix, m_i neutrino mass eigenstates, and ϕ_i relative Majorana CP phases, can be written in terms of oscillation parameters ^{10,11}

$$|m_{ee}^{(1)}| = |U_{e1}|^2 m_1, \quad (3)$$

$$|m_{ee}^{(2)}| = |U_{e2}|^2 \sqrt{\Delta m_{21}^2 + m_1^2}, \quad (4)$$

$$|m_{ee}^{(3)}| = |U_{e3}|^2 \sqrt{\Delta m_{32}^2 + \Delta m_{21}^2 + m_1^2}. \quad (5)$$

The effective mass $\langle m \rangle$ is related with the half-life for $0\nu\beta\beta$ decay via $(T_{1/2}^{0\nu})^{-1} \sim \langle m_\nu \rangle^2$, and for the limit on $T_{1/2}^{0\nu}$ deducible in an experiment we have $T_{1/2}^{0\nu} \sim a \sqrt{\frac{Mt}{\Delta E B}}$. Here a is the isotopical abundance of the $\beta\beta$ emitter; M is the active detector mass; t is the measuring time; ΔE is the energy resolution; B is the background count rate.




Figure 2. Neutrino masses and mixings in the scheme with mass hierarchy. Coloured bars correspond to flavor admixtures in the mass eigenstates ν_1, ν_2, ν_3 . The quantity $\langle m \rangle$ is determined by the dark blue bars denoting the admixture of the electron neutrino U_{ei} .

Neutrino oscillation experiments fix or restrict some of the parameters in (1)–(3), e.g. in the case of normal hierarchy solar neutrino experiments yield Δm_{21}^2 , $|U_{e1}|^2 = \cos^2 \theta_\odot$ and $|U_{e2}|^2 = \sin^2 \theta_\odot$. Atmospheric neutrinos fix Δm_{32}^2 , and experiments like CHOOZ, looking for ν_e disappearance restrict

$|U_{e3}|^2$. The phases ϕ_i and the mass of the lightest neutrino, m_1 are free parameters. Double beta decay can fix the parameter m_1 and thus the absolute mass scale. The expectations for $\langle m \rangle$ from oscillation experiments in different neutrino mass scenarios have been carefully analyzed in ^{10,11}. In sections 2.2.1. to 2.2.3. we give some examples.

2.2.1. Hierarchical Spectrum ($m_1 \ll m_2 \ll m_3$)

In hierarchical spectra (Fig. 2), motivated by analogies with the quark sector and the simplest see-saw models, the main contribution comes from m_2 or m_3 . For the large mixing angle (LMA) MSW solution which is favored at present for the solar neutrino problem (see ³²), the contribution of m_2 becomes dominant in the expression for $\langle m \rangle$, and

$$\langle m \rangle \simeq m_{ee}^{(2)} = \frac{\tan^2 \theta}{1 + \tan^2 \theta} \sqrt{\Delta m_{\odot}^2}. \quad (6)$$

In the region allowed at 90% C.L. by Superkamiokande according to ³³, the prediction for $\langle m \rangle$ becomes

$$\langle m \rangle = (1 \div 3) \cdot 10^{-3} \text{eV}. \quad (7)$$

The prediction extends to $\langle m \rangle = 10^{-2}$ eV in the 99% C.L. range (Fig. 3).

2.2.2. Inverse Hierarchy ($m_3 \approx m_2 \gg m_1$)

In inverse hierarchy scenarios (Fig. 4) the heaviest state with mass m_3 is mainly the electron neutrino, its mass being determined by atmospheric neutrinos, $m_3 \simeq \sqrt{\Delta m_{\text{atm}}^2}$. For the LMA MSW solution one finds ¹¹

$$\langle m \rangle = (1 \div 7) \cdot 10^{-2} \text{eV}. \quad (8)$$

2.2.3. Degenerate Spectrum ($m_1 \simeq m_2 \simeq m_3 \gtrsim 0.1 \text{ eV}$)

In degenerate scenarios (fig. 4) the contribution of m_3 is strongly restricted by CHOOZ. The main contributions come from m_1 and m_2 , depending on their admixture to the electron flavors, which is determined by the solar neutrino solution. We find ¹¹

$$m_{\min} < \langle m \rangle < m_1 \quad \text{with} \quad \langle m_{\min} \rangle = (\cos^2 \theta_{\odot} - \sin^2 \theta_{\odot}) m_1. \quad (9)$$

This leads for the LMA solution to $\langle m \rangle = (0.25 \div 1) \cdot m_1$, the allowed range corresponding to possible values of the unknown Majorana CP-phases.

After these examples we give a summary of our analysis ^{10,11} of the $\langle m \rangle$ allowed by ν oscillation experiments for neutrino mass models in the presently




Figure 3. Double beta decay observable $\langle m \rangle$ and oscillation parameters in the case of the MSW large mixing angle solution of the solar neutrino deficit, where the dominant contribution to $\langle m \rangle$ comes from the second state. Shown are lines of constant $\langle m \rangle$, the lowest line corresponding to $\langle m \rangle = 0.001 \text{ eV}$, the upper line to 0.01 eV . The inner and outer closed line show the regions allowed by present solar neutrino experiments with 90% C.L. and 99% C.L., respectively. Double beta decay with sufficient sensitivity could check the LMA MSW solution. Complementary information could be obtained from the search for a day-night effect and spectral distortions in future solar neutrino experiments as well as a disappearance signal in KAMLAND [from ¹¹].




Figure 4. Left: Neutrino masses and mixings in the inverse hierarchy scenario. Right: Neutrino masses and mixings in the degenerate scheme.

avored scenarios, in Fig. 16. The size of the bars corresponds to the uncertainty in mixing angles and the unknown Majorana CP-phases.




Figure 5. Present sensitivity, and expectation for the future, of the most promising $\beta\beta$ experiments. Given are limits for $\langle m \rangle$, except for the HEIDELBERG-MOSCOW experiment where the recently observed *value*⁸ is given (95% c.l. range), and the best value. Framed parts of the bars: present status; not framed parts: expectation for running experiments; solid and dashed lines: experiments under construction or proposed, respectively. For references see ^{12,85,93}.

2.3 Status of $\beta\beta$ Research and Evidence for the Neutrinoless Decay Mode

The status of present double beta experiments is shown in Fig. 5 and is extensively discussed in ¹². The HEIDELBERG-MOSCOW experiment using the largest source strength of 11 kg of enriched ⁷⁶Ge in form of five HP Ge-detectors is running since August 1990 in the Gran-Sasso underground laboratory ^{34,39,12,1,8}, and is since long time the most sensitive one. We communicate here the status of the analysis of November 2001.

2.3.1. Data from the HEIDELBERG-MOSCOW Experiment

We have analysed ⁸ the data taken in the period August 1990 - May 2000 (54.9813 kg y, or 723.44 mol-years and the data of single site events taken in the period November 1995 - May 2000 with our methods of pulse shape analysis (PSA) ^{19,21,20}, with various statistical methods, in particular with the Bayesian method (see, e.g. ²²). This method is particularly suited for low counting rates, where the data follow a Poisson distribution, that cannot be approximated by a Gaussian.

This Bayesian procedure reproduces γ -lines at the position of known weak

lines from the decay of ^{214}Bi at 2010.7, 2016.7, 2021.8 and 2052.9 keV²⁵. In addition, a line centered at 2039 keV shows up. This is compatible with the Q-value^{27,26} of the double beta decay process. We emphasize, that at this energy no γ -line is expected according to Monte Carlo simulations of our experimental setup, and to the compilations in²⁵. Therefore, on the Bayesian analysis yields a confidence level for a line to exist at 2039.0 keV of 97 % c.l. (2.2σ).

2.3.2. $\beta\beta$ - Half-Life and Effective Neutrino Mass

Under the assumption that the signal at $Q_{\beta\beta}$ is not produced by a background line of at present unknown origin, we can translate the observed number of events into half-lives for neutrinoless double beta decay.

In Table 1 we give the values obtained with the Bayesian method. Also given are the effective neutrino mass $\langle m \rangle$ deduced using matrix elements from²³.

Table 1. Half-life for the neutrinoless decay mode and deduced effective neutrino mass from the HEIDELBERG-MOSCOW experiment⁸.

Significance [kg y]	Detectors	$T_{1/2}^{0\nu}$ y	$\langle m \rangle$ eV	Conf. level
54.9813	1,2,3,4,5	$(0.80 - 35.07) \times 10^{25}$	(0.08 - 0.54)	95% c.l.
54.9813	1,2,3,4,5	$(1.04 - 3.46) \times 10^{25}$	(0.26 - 0.47)	68% c.l.
54.9813	1,2,3,4,5	1.61×10^{25}	0.38	Best Value
46.502	1,2,3,5	$(0.75 - 18.33) \times 10^{25}$	(0.11 - 0.56)	95% c.l.
46.502	1,2,3,5	$(0.98 - 3.05) \times 10^{25}$	(0.28 - 0.49)	68% c.l.
46.502	1,2,3,5	1.50×10^{25}	0.39	Best Value
28.053	2,3,5 SSE	$(0.88 - 22.38) \times 10^{25}$	(0.10 - 0.51)	90% c.l.
28.053	2,3,5 SSE	$(1.07 - 3.69) \times 10^{25}$	(0.25 - 0.47)	68% c.l.
28.053	2,3,5 SSE	1.61×10^{25}	0.38	Best Value

We derive from the data taken with 46.502 kg y the half-life $T_{1/2}^{0\nu} = (0.8 - 18.3) \times 10^{25}$ y (95% c.l.). The analysis of the other data sets, shown in Table 1, and in particular of the single site events data, which play an important role in our conclusion, confirm this result.

The result obtained is consistent with the limits given earlier by the HEIDELBERG-MOSCOW experiment¹. It is also consistent with all other double beta experiments - which still reach less sensitivity. A second Ge-expe-

riment, which has stopped operation in 1999 after reaching a significance of 9 kg y⁵⁷ yields (if one believes their method of 'visual inspection' in their data analysis) in a conservative analysis a limit of $T_{1/2}^{0\nu} > 0.55 \times 10^{25}$ y (90% c.l.). (and not the value given in⁵⁷). The ¹²⁸Te geochemical experiment⁵¹ yields $\langle m_\nu \rangle < 1.1$ eV (68 % c.l.), the ¹³⁰Te cryogenic experiment yields $\langle m_\nu \rangle < 1.8$ eV⁵³, and the CdWO₄ experiment $\langle m_\nu \rangle < 2.6$ eV,⁵² all derived with the matrix elements of²³ to make the results comparable to the present value.

Concluding we obtain, with about 95% probability, first evidence for the neutrinoless double beta decay mode. As a consequence, with 95% confidence, lepton number is not conserved. Further the neutrino is a Majorana particle. The effective mass $\langle m \rangle$ is deduced to be $\langle m \rangle = (0.11 - 0.56)$ eV (95% c.l.), with best value of 0.39 eV. Allowing conservatively for an uncertainty of the nuclear matrix elements of $\pm 50\%$ (for detailed discussions of the status of nuclear matrix elements we refer to¹² and references therein like^{24,18,44}) this range may widen to $\langle m \rangle = (0.05 - 0.84)$ eV (95% c.l.).

In this conclusion, it is assumed that contributions to $0\nu\beta\beta$ decay from processes other than the exchange of a Majorana neutrino (see, e.g.^{12,13} and references therein) are negligible.

2.4 Future of $\beta\beta$ Experiments

To improve the present sensitivity for the effective neutrino mass considerably, and to fix this quantity more accurately, requires new experimental approaches, as discussed extensively in^{12,93,36,28,29,30}. Some of them are indicated in Figs. 5,16.

It has been pointed out earlier (see e.g.^{12,93,36,28,30,36}), that of present generation experiments probably no one has a potential to probe $\langle m \rangle$ below (and perhaps even down to) the present HEIDELBERG-MOSCOW level (see Fig. 5).

The Milano cryogenic experiment using TeO₂ bolometers improved their values for the $\langle m \rangle$ from $\beta\beta$ decay of ¹³⁰Te, from 5.3 eV in 1994⁹⁰ to 1.8 eV in 2000⁵³. NEMO-III, originally aiming at a sensitivity of 0.1 eV, reduced their goals recently to $0.3 \div 0.7$ eV (see⁵⁵) (which is more consistent with estimates given by⁵⁴), to be reached in 6 years from starting of running, foreseen for the year 2002.

2.4.1. GENIUS and other Proposed Future Double Beta Experiments

With the era of the HEIDELBERG-MOSCOW experiment the time of the small smart experiments is over.

To reach significantly larger sensitivity, $\beta\beta$ experiments have to become large. On the other hand source strengths of up to 10 tons of enriched material touch the world production limits. This means that the background has to be reduced by the order a factor of 1000 and more compared to that of the HEIDELBERG-MOSCOW experiment.

Table 2. Some key numbers of future double beta decay experiments (and of the HEIDELBERG-MOSCOW experiment). Explanations: ∇ - assuming the background of the present pilot project. ** - with matrix element from ²³, ²⁴, ⁴⁵, ⁴⁶, ⁴⁷ (see Table II in ³⁵) and assuming an uncertainty of $\pm 50\%$ of the nuclear matrix element. $\triangle$ - this case shown to demonstrate **the ultimate limit** of such experiments. For details see ¹².

$\beta\beta$ - Isotope	Name	Status	Mass (tonnes)	Assumed backgr. $\dagger$ events/ kg y keV, $\ddagger$ events/kg y FWHM, * events /yFWHM	Running Time (tonn. years)	Results limit for $0\nu\beta\beta$ half-life (years)	$\langle m_\nu \rangle$ (eV)
⁷⁶ Ge	HEIDEL- BERG MOSCOW 8,1,41	run- ning since 1990	0.011 (enri- ched)	$\dagger$ 0.06 $\ddagger$ 0.24 * 2	54.98 kg y	(0.8 – 18.3) x 10^{25} y 95% c.l. NOW !!	(0.05-0.84) eV)** 95% c.l. NOW !!
¹⁰⁰ Mo	NEMO III 55	under constr. end 2001?	$\sim$ 0.01 (enri- ched)	$\dagger$ 0.0005 $\ddagger$ 0.2 * 2	50 kg y	10^{24}	0.3-0.7
¹³⁰ Te	CUORE ∇ 91	idea since 1998	0.75 (nat.)	$\dagger$ 0.5 $\ddagger$ 4.5/* 1000	5	$9 \cdot 10^{24}$	0.2-0.5
¹³⁰ Te	CUORE 91,56	idea since 1998	0.75 (nat.)	$\dagger$ 0.005 $\ddagger$ 0.045/* 45	5	$9 \cdot 10^{25}$	0.07-0.2
¹⁰⁰ Mo	MOON 48,59	idea since 1999	10 (enr.) 100(nat.)	?	30 300	?	0.03
¹¹⁶ Cd	CAMEOII CAMEOIII ⁶⁰	idea since 2000	0.65 1(enr.)	* 3. ?	5-8 5-8	10^{26} 10^{27}	0.06 0.02
¹³⁶ Xe	EXO 49,50	Proposal since 1999	1 10	* 0.4 * 0.6	5 10	$8.3 \cdot 10^{26}$ $1.3 \cdot 10^{28}$	0.05-0.14 0.01-0.04
⁷⁶ Ge	GENIUS - TF 42,43	under constr. end 2002	11 kg (enr.)	$\dagger$ $6 \cdot 10^{-3}$	3	$1.6 \cdot 10^{26}$	0.15
⁷⁶ Ge	GENIUS 28,29,30	Pro- posal since 1997	1 (enr.) 1	$\dagger$ $0.04 \cdot 10^{-3}$ $\ddagger$ $0.15 \cdot 10^{-3}$ * 0.15 * 1.5	1 10	$5.8 \cdot 10^{27}$ $2 \cdot 10^{28}$	0.02-0.05 0.01-0.028
⁷⁶ Ge	GENIUS 28,29,30	Pro- posal since 1997	10 (enr.)	$\dagger$ $0.15 \cdot 10^{-3}$ 0 $\triangle$	10 10	$6 \cdot 10^{28}$ $5.7 \cdot 10^{29}$	0.006 - 0.016 0.002 - 0.0056

Table 2 lists some key numbers for GENIUS^{28,29,36}, which was the first proposal for a third generation double beta experiment, and of some other proposals made after the GENIUS proposal. The potential of some of them is shown also in Fig. 16, and it is seen that not all of them will lead to large improvements in sensitivity. Among the latter is also the recently presented MAJORANA project⁵⁸, which does not really apply a striking new strategy for background reduction, particularly also after it was found that the projected segmentation of detectors may not work (see Table 3).

Table 3. Some of the new projects under discussion for future double beta decay experiments (see ref.¹²).

NEW PROJECTS				
	BACKGROUND REDUCTION	MASS INCREASE	POTENTIAL FOR DARK MATTER	POTENTIAL FOR SOLAR ν s
GENIUS	+	+	+	+
CUORE	(+)	+	—	—
MOON	(+)	+	—	+
EXO	+	+	—	—
MAJORANA	—	+	—	—
*) real time measurement of pp neutrinos with threshold of 20 keV (!!)				

For more recent information on XMASS, EXO, MOON experiments see the contributions of Y. Suzuki, G. Gratta and H. Ejiri in Ref. ⁵⁹. The CAMEO project ⁶⁰ in its now propagated variant GEM is nothing then a variant of GENIUS (see below) put into the BOREXINO tank, at some later time. CUORE ⁹¹ has, with the complexity of cryogenic techniques, still to overcome serious problems of background to enter into interesting regions of $\langle m_\nu \rangle$. EXO ⁴⁹ needs still very extensive research and development to probe the applicability of the proposed detection method. In particular if it would be confirmed that tracks will be too short to be identified, it would act essentially only as a highly complicated calorimeter. In the GENIUS project a reduction by a factor of more than 1000 down to a background level of 0.1 events/tonne y keV in the range of $0\nu\beta\beta$ decay is planned to be reached by removing all material close to the detectors, and by using naked Germanium detectors in a large tank of liquid nitrogen. It has been shown that the detectors show excellent performance under such conditions ^{29,38}. For technical questions and extensive Monte Carlo simulations of the GENIUS project for its application in double beta decay we refer to ^{29,38}.

2.4.2. GENIUS and Other Beyond Standard Model Physics

GENIUS will allow besides the large increase in sensitivity for double beta decay described above, the access to a broad range of other beyond SM physics topics in the multi-TeV range. Already now $\beta\beta$ decay probes the TeV scale on which new physics should manifest itself (see, e.g.^{12,28,40,41,37}). Basing to a large extent on the theoretical work of the Heidelberg group in the last six years, the HEIDELBERG-MOSCOW experiment yields results for SUSY models (R-parity breaking, neutrino mass), leptoquarks (leptoquarks-Higgs coupling), compositeness, right-handed W mass, nonconservation of Lorentz invariance and equivalence principle, mass of a heavy left or righthanded neutrino, competitive to corresponding results from high-energy accelerators like TEVATRON and HERA (for details see^{12,40,41}).

2.4.3. GENIUS-Test Facility

Construction of a test facility for GENIUS – GENIUS-TF – consisting of ~ 40 kg of HP Ge detectors suspended in a liquid nitrogen box has been started. Up to summer of 2001, six detectors each of ~ 2.5 kg and with a threshold as low as ~ 500 eV have been produced.

Besides test of various parameters of the GENIUS project, the test facility would allow, with the projected background of 2–4 events/(kg y keV) in the low-energy range, to probe the DAMA evidence for dark matter by the seasonal modulation signature, (see^{42,43}, and section 3).

3 Dark Matter

3.1 SUSY Expectations for Cold Dark Matter

Direct search for WIMPs can be done

(a) by looking for the recoil nuclei in WIMP- nucleus elastic scattering. The signal could be ionisation, phonons or light produced by the recoiling nucleus. The typical recoil energy is a few 100 eV/GeV WIMP mass.

(b) by looking for the modulation of the WIMP signal resulting from the seasonal variation of the earth's velocity against the WIMP 'wind'.

The expectation for neutralino elastic scattering cross sections and masses have been extensively analysed in many variants of SUSY models.

Figs.6,7 represent the present situation. The SUSY predictions in Figs.6a,7 are from the MSSM with relaxed unification conditions⁶⁶ and the MSUGRA model⁶⁷. Fig.6b shows the result of a study 'at Post-LEP Benchmark points' based again on the MSUGRA⁶⁸. Present experiments only just touch the border of the area predicted by the MSSM. The experimental DAMA evidence for dark matter lies in an area, in which MSUGRA mod-




Figure 6. (a): WIMP-nucleon cross section limits in pb for scalar interactions as function of the WIMP mass in GeV. Shown are contour lines of present experimental limits (solid lines) and of projected experiments (dashed lines). Also shown is the region of evidence published by DAMA. The theoretical expectations from the MSSM are shown by two scatter plots, - for accelerating and for non-accelerating Universe (from⁶⁶), and from the SUGRA by the grey region (from⁶⁷). Only GENIUS will be able to probe the shown range also by the signature from seasonal modulations. (b): WIMP- proton elastic scattering cross sections according to various MSUGRA models (see text). From⁶⁸.

els do not expect dark matter. They would require beyond GUT physics in this frame⁶⁹.

It is clear from extensive theoretical work that high-sensitivity dark matter experiments can yield an important contribution to SUSY search. Fig.8 (from⁷⁰) shows, in the MSUGRA model, the SUSY reach contours for different accelerators (LEP2, Tevatron, LHC, NLC) together with direct detection rates in a ^{73}Ge detector. It is visible that a detector of GENIUS-sensitivity operates in SUSY search on the level of LHC and NLC. Fig.9 (from⁶⁶), shows another study in the MSSM with relaxed unification conditions. Non-observation of Dark Matter with GENIUS would exclude a 'light' SUSY spectrum (all sfermion masses lighter than 300 - 400 GeV) and any possibility for a light Higgs sector in the MSSM.

If classifying the SUGRA models into more $g_{\mu-2}$ -friendly (I,L,B,G,C,J) and less $g_{\mu-2}$ -friendly models, according to⁶⁸, the former ones have good prospects to be detectable by LHC and/or a 1 TeV collider. GENIUS could check not only the larger part of these ones, but in addition two of the less $g_{\mu-2}$ -friendly models (E and F), which will be difficult to be probed by fu-




Figure 7. The same in Figs.6a, but for spin-dependent interaction. The grey area corresponds to the MSUGRA expectations⁶⁷. Full lines denote present experimental results, dashed lines expectations of future experiments.

ture colliders (see Fig.6b). This demonstrates nicely the complementarity of collider and underground research.

It might be mentioned that in case of $g_{\mu-2}$ - unfriendly models, i.e. those with very low cross sections in Figs.6b, it might be required to turn from spin-zero targets and looking for spin-independent interaction, which usually for not too light nuclei gives the largest cross sections, to spin- non-zero target nuclei and spin-dependent interaction⁶⁶. It has been shown recently⁶⁶ that if spin-zero experiments with sensitivities of 10^{-5} - 10^{-6} events/kg day will fail to detect a dark matter signal, an experiment with nonzero spin target and higher sensitivity will be able to detect dark matter *only* due to the spin neutralino-quark interaction (see Fig.10).

3.2 Cold Dark Matter - Present and Future Experiments

Summarizing the present experimental status, present and also future projects can be categorized in two classes:

1. Sensitivity (or sensitivity goal) 'just for' confirmation of DAMA.
2. Sensitivity to enter deeply into the range of SUSY predictions.

Only very few experiments may become candidates for category 2 in a foreseeable future (see Figs.6,7), and as far as at present visible, of those only




Figure 8. Direct detection rate R in events/kg day in a ^{73}Ge detector, and SUSY reach contours for LEP2, Tevatron (MI), LHC slepton signal and NLC. TH - excluded by theoretical considerations. EX - excluded by collider searches for SUSY particles (from⁷⁰).

GENIUS will have the chance to search for modulation, i.e. to check, like DAMA, positive evidence for a dark matter signal.

Figs.11,12 give an overview of present and future experiments. The at present most sensitive experiments DAMA, CDMS (and Edelweiss) are claimed⁷¹ not to be fully consistent, although CDMS can at present not exclude the full DAMA evidence region^{73,71}. Some problems in the data analysis of CDMS have been revised recently⁷².

One of the main problem of the cryodetectors is to obtain good numbers of background in the *raw* data, i.e. of the starting values for the rejection procedure. This is also the reason, why simple cryogenic calorimeters such as the CRESST I^{82,74} and the Milano^{53,89} bolometers are not fully competitive.

The at present most sensitive experiments using raw data are the HEIDELBERG-MOSCOW experiment⁸⁰ and the HEIDELBERG Dark Matter Search Experiment (HDMS)^{83,81}. HDMS uses a special configuration of Ge detectors, with an enriched ^{73}Ge detector embedded into a natural Ge-detector (Fig.13a). Results of the pilote project are given in^{83,81}. The final configuration with the enriched ^{73}Ge inner detector is in operation since




Figure 9. Different lower bounds for the total event rate in ^{73}Ge (events/day kg) versus mass of the LSP (GeV). Here $M_{sq, CH, H}$ denote masses of the squark, the charged Higgs boson, and the light neutral CP-even Higgs boson, respectively. Heavy chargino mass is denoted as M_{ch-os} . “Full” corresponds to the lower bound obtained from main (unconstrained) scan, and “light spectrum” denotes the lower bound for R, which is obtained with all sfermion masses lighter than about 300 GeV. The horizontal dotted line represents expected sensitivity for the direct dark matter detection with GENIUS (from⁶⁶).

February 2001.

The experimental future is also illustrated in Figs.6,7. A useful overview is given in⁶⁴, and for Ge detectors in⁷⁹. For an earlier review see⁸⁷.

The cryogenic projects are CDMS II, CREST II, Edelweiss II. In contrast to CDMS and Edelweiss which do its active background rejection by looking for ionisation and phonons, CRESST II plans to use simultaneous detection of light and phonons. CDMS II plans to use 42 detectors with a total mass of 6.8 kg of Ge by 2006 in the Soudan mine, CRESST II plans to have ~ 10 kg of CdWO_4 in the Gran Sasso in some future. The cryogenic experiments are, however, operating at present only 600 g of detectors *or less*, after a decade of development. CDMS has collected only 10.6 kg d of data over this time (in 1999, since then *no* measurement)⁷¹, Edelweiss only 4.53 kg d⁷⁵. Therefore, they may have severe problems to expand their small detector masses to several tens of kg or better 100 kg, as required for modulation search. This means that although e.g. CDMS II may reach a future sensitivity in an exclusion plot as shown in Fig.6b, it will not be able to look for the modulation signal. A general problem in the present stage still seems to be the reproducibility of the highly complicated cryo detectors. In spite of this, phantasy is large enough, to dream already about 1 ton cryo detectors systems⁷¹.




Figure 10. Ratio of spin-dependent event rate to spin-independent event rate in ^{73}Ge as function of LSP mass (upper left), of total (spin-dependent plus spin-independent) event rate (lower left part), and of scalar cross section of neutralino-proton interaction (lower right) obtained with $0.1 < \Omega_\chi h_0^2 < 0.3$. The vertical line gives the expected sensitivity of GENIUS. In the region above the horizontal line the spin contribution dominates. The total event rate versus gaugino fraction of LSP is also given (upper right). From⁶⁶.

Other far future projects are the superheavy droplet detectors PICASSO/SIMPLE⁶⁴. They are working at present on a scale of 15 and 50 g detectors. Their idea is to use 10-100 μm diameter droplets of volatile C_4F_{10} , C_3F_8 , ... in metastable superheated condition and to choose critical energy and radius such that only nuclear recoils can trigger a phase transition, but not γ and β particles. The acoustic signal of the explosive bubble formation will be observed. The expected sensitivity of a 1 ton module for spin-dependent WIMP-nucleon interaction is shown in Figure 6b - as SDD 1 ton (for an assumed U/Th contamination of 10^{-15} g/g - U/Th α -emitters can cause recoil events!). A drawback is that these detectors cannot measure energy spectra of WIMPs.

A very promising project which would yield a nice signal identification, is DRIFT in the Boulby mine. It is aiming at looking for the diurnal directional

The Potencial of World Dark Matter Underground Experiments				
NAME	TARGET	MASS (kg)	MODULATIONS	UPPER LIMIT
DAMA-I	NaI(Tl)	100.	YES 2 - 20 keV 57986.0 kg day ⁺	Runs since 1996
DAMA-II	NaI(Tl)	250.	YES	Plans for 2002
HDMS	⁷³ Ge	0.2	NO 0.07 ev/kg d keV 40 - 100 keV 0.2 ev/kg d keV 11 - 40 keV	Runs since 2001
GENIUS-TF	⁷⁶ Ge	40.	YES 6 new det. 2.5 kg threshold 500 eV !!!	In construction start in 2002
GENIUS	⁷⁶ Ge	100. 1000.	YES YES	Proposal since 1997 Proposal since 1997
CDMS-I	Ge (blip)	0.495 (4 x 0.163) 3 (interm, 1 anticoin.)	NO 10-100 keV 50(int) - 230(ext.) kg d ⁺	Run 96 live days in period: 11.98-09.99
CDMS-II	Ge (blip)	10.6	NO	Plan for 2006
CRESST-I	CaWO ₄	1.048	NO 1.2 c/keV kg d at 15 keV	Run in 1998 some months
CRESST-II	CaWO ₄	10.	NO	Plans since 2000

Figure 11. Status of some running and future Dark Matter experiments.

Direct detection:

- Ionization detectors - Ge
HEIDELBERG-MOSCOW
HDMS (Heidelberg Dark Matter Search exp.
IGEX
-Scintillation detectors - NaI
DAMA NAIAD
- Cryodetectors
CDMS
EDELWEISS
CRESST
LiF bolometers
ROSEBUD (CAVENDISH)
- Xe detector
ZEPLIN II
- Superheated droplet
PICASSO / SIMPLE
- Drift chamber (diurnal modulation)
DRIFT
- New Ionization Detector Approaches
GENIUS - TF
GENIUS GEDEON

Indirect detection:

- Ice Cerenkov Detector (AMANDA)
- Water Cerenkov Detector (ANTARES)

Figure 12. List of some of the main present and future dark matter experiments.

modulation (Fig.13b). The idea is to detect tracks of nuclear recoils in a TPC with Xe(Ar) by a multiwire read-out. A 1 m^3 prototype (1.5 kg Xe) is under construction. It is seen as first component of the full 10 m^3 DRIFT experiment^{64,65}.

The ZEPLIN project uses scintillation and electro-luminescence in two-phase xenon. Plans for ZEPLIN II are 0.01 - 0.1 counts/kg d and 20 kg of Xe. While still waiting for results of ZEPLIN I, plans are already discussed for a ZEPLIN IV⁷⁶.

To return to the more 'earth-bound' projects: DAMA will extend their mass to 250 kg, and plans to start operation in summer 2002⁷³. Also the NAIAD project (Boulby mine) plans to use NaI - 40-100 kg of NaI in a liquid scintillator Compton veto⁷⁷. The projected NAIAD limits for 100 kg y exposure are 0.1 c/kg d. Because of the large mass it will be able to look for modulation.

The HDMS experiment and the GENIUS-TF experiment^{42,43} aim at probing the DAMA evidence (see Fig.15). GENIUS-TF consisting of 40 kg of Ge detectors in liquid nitrogen (Fig.14a) could also measure the modulation signal^{43,78}. Up to summer 2001, already 6 detectors of 2.5 kg each, with an extreme low-energy threshold of $\sim 500\text{ eV}$ have been produced. A similar potential is aimed at by the GEDEON project⁷⁹, which plans to use 28 Ge diodes in one single cryostat.

GENIUS-TF is already under installation in the Gran-Sasso laboratory and should start operation by end of 2002.

The probably most far reaching project is GENIUS^{12,29}. Since it is based on conventional techniques, using Ge detectors in liquid nitrogen, it may be realized in the most straightforward way.

GENIUS would already in a first step, with 100 kg of *natural* Ge detectors in three years of measurement, cover a significant part of the SUSY parameter space for prediction of neutralinos as cold dark matter (Figs. 6,7). For this purpose the background in the energy range $< 100\text{ keV}$ has to be reduced to 10^{-2} (events/kg y keV). At this level solar neutrinos as source of background are still negligible. Of particular importance is to shield the detectors during production (and transport) to keep the background from spallation by cosmic rays sufficiently low (for details see^{36,85,88}).

The sensitivity of GENIUS for Dark Matter corresponds to that obtainable with a 1 km^3 AMANDA detector for *indirect* detection (neutrinos from annihilation of neutralinos captured at the Sun) (see⁹²). Interestingly both experiments would probe different neutralino compositions: GENIUS mainly gaugino-dominated neutralinos, AMANDA mainly neutralinos with comparable gaugino and Higgsino components (see Fig. 38 in⁹²).




Figure 13. HDMS and DRIFT experiments.

3.3 Hot Dark Matter Search

According to the recent indication for the neutrinoless mode of double beta decay⁸, (see section 2), neutrinos should still play an important role as hot dark matter in the Universe.

The effective mass has been determined to be⁸ $\langle m \rangle = (0.05 - 0.84) \text{ eV}$ at a 95% c.l. (best value 0.39 eV) *including* an uncertainty of $\pm 50\%$ of the nuclear matrix elements.

With the limit deduced for the effective neutrino mass, the HEIDELBERG-MOSCOW experiment excludes several of the neutrino mass scenarios allowed from present neutrino oscillation experiments (see Fig.16) - allowing mainly only for degenerate and partially degenerate mass scenarios and an inverse hierarchy 3ν - scenario (the latter being, however, strongly disfavored by a recent analysis of SN1987A). In particular hierarchical mass schemes are excluded.

Assuming the degenerate scenarios to be realized in nature we fix - according to the formulae derived in¹⁰ - the common mass eigenvalue of the




Figure 14. (a): Conceptual design of the Genius TF. Up to 14 detectors will be housed in the inner detector chamber, filled with liquid nitrogen. As a first shield 5 cm of zone refined Germanium, or extremely low-level copper will be used. Behind the 20 cm of polystyrene isolation another 35 cm of low level lead and a 15 cm borated polyethylene shield will complete the setup. (b): GENIUS - 100 kg of Ge detectors are suspended in a large liquid nitrogen tank (see^{37,29,12}).




Figure 15. The potential of GENIUS-TF for dark matter search. WIMP-nucleon cross section limits as a function of the WIMP mass for spin-independent interactions. The solid lines are current limits of the HEIDELBERG-MOSCOW experiment, the HDMS, the DAMA and the CDMS experiments. The dashed curves are the expectation for HDMS, and for Genius-TF with an energy threshold of 11 keV and 2 keV (no tritium contamination) respectively, and a background index of 2 events/kg y keV below 50 keV. The filled contour represents the evidence region of the DAMA experiment




Figure 16. The impact of the evidence obtained for neutrinoless double beta decay⁸ (best value of the effective neutrino mass $\langle m \rangle = 0.39$ eV, 95% confidence range (0.05 - 0.84) eV - allowing already for an uncertainty of the nuclear matrix element of a factor of $\pm 50\%$, on possible neutrino mass schemes. The bars denote allowed ranges of $\langle m \rangle$ in different neutrino mass scenarios, still allowed by neutrino oscillation experiments. Hierarchical models are excluded by the new $0\nu\beta\beta$ decay result. Also shown are the expected sensitivities for the future potential double beta experiments CUORE, MOON, EXO, and the 1 ton and 10 ton project of GENIUS^{29,36,88,85} (from⁹).

degenerate neutrinos to $m = (0.05 - 3.4)$ eV. Part of the upper range is already excluded by tritium experiments, which give a limit of $m < 2.2$ eV (95% c.l.)⁹⁴. The full range can only partly (down to ~ 0.5 eV) be checked by future tritium decay experiments, but could be checked by some future $\beta\beta$ experiments (see, e.g.^{12,88,85}). The deduced best value for the mass is consistent with expectations from experimental $\mu \rightarrow e\gamma$ branching limits in models assuming the generating mechanism for the neutrino mass to be also responsible for the recent indication for an anomalous magnetic moment of the muon⁶¹. It lies in a range of interest also for Z-burst models recently discussed as explanation for super-high energy cosmic ray events beyond the GKZ-cut-off⁸⁴. The sensitivity of the present result is already in the range to be probed by the satellite experiments MAP and PLANCK (Fig.17).

4 Conclusion

Dark matter search is presently one of the most exciting fields of particle physics and cosmology. Underground experiments at present only marginally

$m_1(\text{eV})$ HEIDELBERG-MOSCOW Positive EVIDENCE




Figure 17. Double beta decay observable $\langle m \rangle$ and oscillation parameters: The case for degenerate neutrinos. Plotted on the axes are the overall scale of neutrino masses m_0 and mixing $\tan^2 \theta_{12}$. Also shown is a cosmological bound deduced from a fit of CMB and large scale structure³¹ and the expected sensitivity of the satellite experiments MAP and PLANCK. The present limit from tritium β decay of 2.2 eV^{94} would lie near the top of the figure. The range of $\langle m \rangle$ fixed by the HEIDELBERG-MOSCOW experiment is, in the case of small solar neutrino mixing, already in the range to be explored by MAP and PLANCK.

touch in their sensitivity the range of present SUSY predictions for cold dark matter. Of future experiments the GENIUS project has the best prospects to cover a large part of the predicted range. GENIUS will provide information complementary to future collider search. This information is indispensable, even if LHC would find supersymmetry, since in any case it still has to be shown that SUSY particles indeed form the cold dark matter in the Universe. GENIUS will simultaneously be the most straightforward way to fix the neutrino mass and the contributions of neutrinos to hot dark matter with higher accuracy.

References

1. H.V. Klapdor-Kleingrothaus et al.,(HEIDELBERG-MOSCOW Collaboration), *Eur. Phys. J A* **12** (2001) 147 - 154 and hep-ph/0103062 and in Proc. of DARK2000, H.V. Klapdor-Kleingrothaus (Ed.), Springer-Verlag, Heidelberg, (2001) 520-533 and [http : // www.mpi - hd.mpg.de / non_acc /](http://www.mpi-hd.mpg.de/non_acc/).
2. E. Majorana, *Nuovo Cimento* **14** (1937) 171 - 184.

3. G. Racah, *Nuovo Cimento* **14** (1937) 322 - 328.
4. W.H. Furry, *Phys. Rev.* **56** (1939) 1184 - 1193.
5. J. A. McLennan, Jr. *Phys. Rev.* **106** (1957) 821.
6. K. M. Case, *Phys. Rev.* **107** (1957) 307.
7. D. V. Ahluwalia, *Int. J. Mod. Phys. A* **11** (1996) 1855.
8. H.V. Klapdor-Kleingrothaus, A. Dietz, H.L. Harney and I.V. Krivosheina; *Modern Physics Letters A* **16**, No. **37** (2001) 2409-2420.
9. H.V. Klapdor-Kleingrothaus and U. Sarkar, *Modern Physics Letters A* **16**, No. **38** (2001) 2469-2482.
10. H.V. Klapdor-Kleingrothaus, H. Päs and A.Yu. Smirnov, *Phys. Rev. D* **63** (2001) 073005 and *hep-ph/0003219*.
11. H.V. Klapdor-Kleingrothaus, H. Päs and A.Yu. Smirnov, in *Proc. of DARK'-2000*, Heidelberg, 10-15 July, 2000, Germany, ed. H.V. Klapdor-Kleingrothaus, *Springer, Heidelberg* (2001) 420 - 434.
12. H.V. Klapdor-Kleingrothaus, "60 Years of Double Beta Decay - From Nuclear Physics to Beyond the Standard Model", *World Scientific, Singapore* (2001) 1281 p.
13. R.N. Mohapatra and P.B. Pal, "Massive Neutrinos in Physics and Astrophysics", *Singapore, Singapore: World Scientific, World Scientific lecture notes in physics*, **41** (1991) 318 pp.
14. M. Doi et al., *Prog. of Theor. Phys. Suppl.* **83** (1985) 1-175.
15. K. Muto and H.V. Klapdor, in "Neutrinos", *Graduate Texts in Contemporary Physics*, ed. H.V. Klapdor, Berlin, Germany: Springer (1988) 183 - 238.
16. K. Grotz and H.V. Klapdor, "Die Schwache Wechselwirkung in Kern-, Teilchen- und Astrophysik", B.G. Teubner, Stuttgart (1989), "The Weak Interaction in Nuclear, Particle and Astrophysics", IOP Bristol (1990), Moscow, MIR (1992) and China (1998).
17. H.V. Klapdor-Kleingrothaus and A. Staudt, "Teilchenphysik ohne Beschleuniger", B.G. Teubner, Stuttgart (1995), "Non-Accelerator Particle Physics", IOP Publishing, Bristol and Philadelphia (1995) and 2. ed. (1998) and Moscow, Nauka, Fizmatlit (1998), translated by V.A. Bednyakov.
18. P.Vogel in "Current Aspects of Neutrino Physics", ed. D.O. Caldwell, Berlin, Heidelberg, Germany: Springer (2001) 177 - 198.
19. J. Hellmig and H.V. Klapdor-Kleingrothaus, *Nucl. Instrum. Meth. A* **455** (2000) 638 - 644.
20. J. Hellmig, F. Petry and H.V. Klapdor-Kleingrothaus, *Patent* **DE19721323A**.
21. B. Majorovits and H.V. Klapdor-Kleingrothaus. *Eur.Phys.J.* **A6** (1999) 463.
22. G. D'Agostini, hep-ex/0002055, W. von der Linden and V. Dose, *Phys. Rev. E* **59** 6527 (1999), and F.H. Fröhner, JEFF Report **18** NEA OECD (2000) and *Nucl. Sci. a. Engineering* **126** (1997) 1, K. Weise and W. Wöger, "Messunsicherheit und Messdatenauswertung", Wiley-VCH, Weinheim 1999, and P.M. Lee, "Bayesian Statistics: An Introduction", Second edition, Arnold, London 1997, and A. O'Hagan, "Bayesian Inference", Volume 2B of "Kendall's Ad-

- vanced Theory of Statistics", Arnold, London, 1997.
23. A.Staudt, K.Muto and H.V.Klapdor-Kleingrothaus, *Eur. Lett.* **13** (1990) 31.
 24. T. Tomoda, *Rept. Prog. Phys.* **54** (1991) 53 - 126.
 25. R.B. Firestone and V.S. Shirley, Table of Isotopes, 8-th Edition, *John Wiley and Sons*, Incorp., N.Y. (1998).
 26. G. Douysset et al., *Phys. Rev. Lett.* **86** (2001) 4259 - 4262.
 27. J.G. Hykawy et al., *Phys. Rev. Lett.* **67** (1991) 1708.
 28. H.V. Klapdor-Kleingrothaus in Proc. of BEYOND'97, Castle Ringberg, Germany, 8-14 June 1997, eds. by H.V. Klapdor-Kleingrothaus and H. Päs, *IOP Bristol* (1998) 485 - 531.
 29. H.V. Klapdor-Kleingrothaus et al. **MPI-Report MPI-H-V26-1999** and *Preprint: hep-ph/9910205* and in Proc. of BEYOND'99, Castle Ringberg, Germany, 6-12 June 1999, edited by H.V. Klapdor-Kleingrothaus and I.V. Krivosheina, *IOP Bristol* (2000) 915 - 1014.
 30. H.V. Klapdor-Kleingrothaus, in Proc. of NEUTRINO 98, Takayama, Japan, 4-9 Jun 1998, (eds) Y. Suzuki et al. *Nucl. Phys. Proc. Suppl.* **77** (1999) 357.
 31. R.E. Lopez, *astro-ph/9909414*; J.R. Primack and M.A.K. Gross, *astro-ph/0007165*; J.R. Primack, *astro-ph/0007187*; J. Einasto, in Proc. of DARK2000, Heidelberg, Germany, July 10-15, 2000, Ed. H.V. Klapdor-Kleingrothaus, *Springer, Heidelberg* (2001) 3-11.
 32. Y. Suzuki in Proc. of NEUTRINO2000, ed. J. Law et al. (2001), *Nucl. Phys. B* **91** (2001) 29 - 35.
 33. M.C. Gonzalez-Garcia et al., *Phys. Rev. D* **63** (2001) 033005.
 34. HEIDELBERG-MOSCOW Coll. (M. Günther et al.), *Phys.Rev.D* **55** (1997) 54.
 35. HEIDELBERG-MOSCOW Coll., *Phys. Rev. Lett.* **83** (1999) 41 - 44.
 36. H.V. Klapdor-Kleingrothaus, in Proc. of Int. Conference NOW2000 - "Origins of Neutrino Oscillations", ed. G. Fogli, *Nucl. Phys. B* **100** (2001) 309 - 313.
 37. H.V. Klapdor-Kleingrothaus, *Int. J. Mod. Phys. A* **13** (1998) 3953.
 38. H.V.Klapdor-Kleingrothaus, J.Hellmig, M.Hirsch, *J. Phys. G* **24** (1998) 483.
 39. H.V. Klapdor-Kleingrothaus, in Proc. of Advances in Nuclear Physics, eds.: D. Poenaru and S. Stoica, *World Scientific, Singapore* (2000) 123 - 129.
 40. H.V. Klapdor-Kleingrothaus, in Proc. of 'Lepton-Baryon98', Trento, Italy, 20-25 April, 1998, ed. H.V. Klapdor-Kleingrothaus and I.V. Krivosheina, *IOP, Bristol*, (1999) 251-301 and *Preprint: hep-ex/9901021*.
 41. H.V. Klapdor-Kleingrothaus, *Springer Tracts in Modern Physics*, **163** (2000) 69-104, *Springer-Verlag, Heidelberg, Germany* (2000).
 42. H.V. Klapdor-Kleingrothaus, L.Baudis, A.Dietz, G.Heusser, I.Krivosheina, B.Majorovits and H. Strecker, Internal Report **MPI-H-V32-2000**.
 43. H.V. Klapdor-Kleingrothaus et al., *hep-ph/0103082*, subm. for publ. (2001).
 44. A. Faessler and F. Simkovic, *J. Phys. G* **24** (1998) 2139 - 2178.
 45. W.C. Haxton and G.J. Stephenson, *Prog. Part. Nucl. Phys.* **12** (1984) 409.
 46. X.R. Wu, A. Staudt, H.V. Klapdor-Kleingrothaus, Cheng-Rui Ching and Tso-Hsiu Ho, *Phys. Lett. B* **272** (1991) 169 - 172.

47. X.R. Wu, A. Staudt, T.T.S. Kuo and H.V. Klapdor-Kleingrothaus, *Phys. Lett. B* **276** (1992) 274 - 278.
48. H. Ejiri et al., *Phys. Rev. Lett.* **85** (2000) 2917-2920.
49. M. Danilov et al., *Phys. Lett. B* **480** (2000) 12 - 18.
50. G. Gratta in Proc. of LowNu2, Dec. 4-5 (2000) Tokyo, Japan, ed: Y. Suzuki, World Scientific (2001).
51. T. Bernatowicz et al., *Phys. Rev. Lett.* **69** (1992) 2341 - 2344
52. F.A. Danevich et al., *Phys. Rev. C* **62** (2000) 045501.
53. A. Alessandrello et al., *Phys. Lett. B* **486** (2000) 13-21 and S. Pirro et al., *Nucl. Instr. Methods A* **444** (2000) 71-76.
54. V.I. Tretyak and Yu.G. Zdesenko, *At. Data Nucl. Data Tables* **61** (1995) 43.
55. NEMO Coll., Contr. paper for XIX Int. Conf. NEUTRINO2000, Sudbury, Canada, June 16 - 21, 2000 LAL 00-31 (2000) 1 - 10 and in Proc. of NAN-Pino2000, Dubna, Russia, July 2000, ed. V. Bednjakov et al. (2001) 1296.
56. E. Fiorini in Proc. of NEUTRINO2000, ed. J. Law et al. *Nucl. Phys. B* **91** (Proc. Suppl.) (2001) 262 - 269.
57. C.E. Aalseth et al. (IGEX Collaboration), *Yad. Fiz.* **63**, No 7 (2000) 1299.
58. L. DeBraekeleer, talk at Work.. on the Next Gener. U.S. Undergr. Sci. Fac., WIPP, June 12-14, 2000, Carlsbad, New Mexico, USA and C.E. Aalseth et al., in Proc. of TAUP'2001, Gran Sasso, Italy September 2001 ed. A. Bettini.
59. Proc. Int. Workshop on Low Energy Solar Neutrinos, LowNu2, December 4 and 5 (2000) Tokyo, Japan, ed: Y. Suzuki, World Scientific, Singapore (2001).
60. G. Bellini et al. *Phys. Lett. B* **493** 216 - 228 (2000).
61. E. Ma and M. Raidal, *Phys. Rev. Lett.* **87** (2001) 011802; Erratum-ibid. **87** (2001) 159901.
62. K. Freese et al. in Proc. of DARK'98, Heidelberg, July 1998, *IOP, Bristol* (1999) eds. *H.V. Klapdor-Kleingrothaus et al.* 352; D. Graff, in Proc. of DARK 2000, Heidelberg, July 2000, Springer, Heidelberg (2001) ed. *H.V. Klapdor-Kleingrothaus*, 352 and K. Freese et al. in Proc. of DM'2000, WS (2001) 213 - 223 and *astro-ph/0007444*.
63. S. Asztalos et al., *Phys. Rev. D* **64** (2001) 092003; G.G. Raffelt in Proc. of DARK'2000, Heidelberg, July 2000, *Springer, Heidelberg* (2001), ed. *H.V. Klapdor-Kleingrothaus*, 60.
64. V. Zacek NPB, in Proc. of NEUTRINO2000, *Sudbury, Canada*, June 2000, ed. *J. Law et al.* (2001), *Nucl. Phys. Proc. Suppl.* **91** (2001) 368.
65. M.J. Lehner et al., in Proc. of DARK'2000, Heidelberg, July 2000, *Springer, Heidelberg* (2001), ed. *H.V. Klapdor-Kleingrothaus*, 590-597, C.J. Martoff, DRIFT Collab. in Proc. York 2000, ed. *J.C. Spooner et al.* 463-474.
66. V.A. Bednyakov and H.V. Klapdor-Kleingrothaus, *Phys. Rev. D* **62** (2000) 043524/1, and *Phys. Rev. D* **63** (2001) 095005.
67. J. Ellis et al. *Phys.Lett. B* **481** (2000) 304 and *Phys.Rev. D* **63** (2000) 065016.
68. J. Ellis et al., *hep-ph/0111294*.
69. R. Arnowitt, private communication, 2001

70. M. Brhlik in Proc. of DARK'98, Heidelberg, July 1998, IOP, Bristol (1999) eds. *H.V. Klapdor-Kleingrothaus et al.* 499-515.
71. R.J. Gaitskell, NANP2001, Dubna, Russia, June 2001, to be publ. in *Nuclei, Letters* (2002).
72. B. Sadoulet in Proc.TAUP'2001, Sept.2001, Gran Sasso, Italy, ed. *A. Bettini*.
73. R. Bernabei et al., in Proc.: of Erice'2001, Sept.2001, Erice, Italy, ed. *A. Faessler* and of TAUP'2001, Sept.2001, Gran Sasso, Italy, ed. *A. Bettini*.
74. F. Pröbst in Proc. of TAUP'2001, Sept. 2001, Gran Sasso, Italy, ed. *A. Bettini*.
75. J. Gascon et al. in Proc.of DARK'2000, Heidelberg, July 2000, *Springer, Heidelberg* (2001), ed. *H.V. Klapdor-Kleingrothaus*, 575-580 and A. de Lesquen et al. in Proc. of NANP2001, Dubna, Russia, June 2001, ed. *V. Bednjakov*, to be publ. in *Nuclei, Letters* (2002).
76. D.B. Cline, *astro-ph/0111098*.
77. J.C. Spooner in Proc. of York'2000, ed. *J.C. Spooner et al.* WS (2001).
78. H.V. Klapdor-Kleingrothaus et al., subm for publ. 2001.
79. A. Morales., *hep-ex/0111089*.
80. HEIDELBERG-MOSCOW Collaboration, *Phys. Rev. D* **59** (1998) 022001.
81. L. Baudis, A. Dietz, B. Majorovits, F. Schwamm, H. Strecker and H.V. Klapdor-Kleingrothaus, *Phys. Rev. D* **63** (2000) 022001.
82. Y. Ramachers for the CRESST Coll. in Proc. of XIth Renc. de Blois, Front. of Matter, France, June 27-July 3, 1999.
83. H.V. Klapdor-Kleingrothaus et al. in Proc. of DARK2000, Heidelberg, Germany, July 10-15, 2000, *Springer, Heidelberg* (2001), ed. *H.V. Klapdor-Kleingrothaus* 553- 568.
84. T.J. Weiler, in Proc. Beyond the Desert 1999, Tegernsee, Germany, 6-12 Juni 1999, edited by H.V. Klapdor-Kleingrothaus and I.V. Krivosheina, IOP Bristol (2000) 1085 - 1106; H. Päs and T.J. Weiler, *Phys. Rev. D* **63** (2001) 113015.
85. H.V. Klapdor-Kleingrothaus, in Proc. of NOON2000, Tokyo, Dec. 2000, *World Scientific, Singapore* (2001).
86. M. Tegmark, M. Zaldarriaga and A.J.S. Hamilton, *Preprint: hep-ph/0008145*.
87. H.V. Klapdor-Kleingrothaus and Y. Ramachers, *Eur.Phys.J. A* **3** (1998) 85-92.
88. H.V. Klapdor-Kleingrothaus, in Proc. of LowNu2, Dec. 4-5., Tokyo, Japan, ed: *Y. Suzuki et al. World Scientific, Singapore* (2001).
89. E. Fiorini et al., *Phys. Rep.* **307**, 309 (1998).
90. A. Alessandrello et al., *Phys. Lett. B* **335**, 519-525 (1994).
91. A. Giuliani (CUORE Collaboration), in Proc. Astrophysics and Cosmology', 'Lepton-Baryon98',eds. *H.V. Klapdor-Kleingrothaus and I.V.Krivosheina*, Trento, Italy, April 20 - April 25, 1998, *IOP, Bristol*, 302 - 308 (1999).
92. J. Edsjö, home page: <http://www.physto.se/edsjo/>
93. H.V. Klapdor-Kleingrothaus, in Proc. of NANPino-2000, Dubna, July 19-22, 2000, Part. and *Nucl. Lett.* iss. 1/2 (2001) 20-39 and *hep-ph/0102319*.
94. C. Weinheimer, in Proc. of DARK2000, Heidelberg, Germany, 10-16 July, 2000, ed. H.V. Klapdor-Kleingrothaus, *Springer, Heidelberg* (2001) 513 - 519.